



Research
Education
Outreach

CCA

A Nonlinear Sandwich Theorem

Mario Ghossoub, Giulio Principi and Lorenzo Stanca

No. 694

July 2023

Carlo Alberto Notebooks

www.carloalberto.org/research/working-papers

A NONLINEAR SANDWICH THEOREM

MARIO GHOSSOUB
UNIVERSITY OF WATERLOO

GIULIO PRINCIPI
NEW YORK UNIVERSITY

LORENZO STANCA
COLLEGIO CARLO ALBERTO AND UNIVERSITY OF TURIN

APRIL 5, 2023

ABSTRACT. We provide a Sandwich Theorem ([König \(1972\)](#)) for positively homogeneous functionals that satisfy additivity only on a restricted domain. Our relaxation of additivity is based on a binary relation called *convex-conic symmetric preorder*, whereby additivity is restricted to all couples of elements that belong to such relation. We then study applications of our nonlinear Sandwich Theorem, proving extension and envelope representation results. Finally, we consider some applications to comonotonicity, a key property in decision theory, risk measurement, and the theory of risk sharing.

1. INTRODUCTION

The classic version of the Sandwich Theorem yields the existence of a linear functional in between a given sublinear functional and a given superlinear functional. [König \(1972\)](#) proved this result as a corollary to the Hahn-Banach extension theorem. In a recent paper, [Amarante \(2019\)](#) provided an alternative proof, by combining an argument in [Fuchssteiner and Wright \(1977\)](#) with Pataia's Fixed Point Theorem ([Pataia \(1997\)](#)). In this paper, we show how the elegant approach developed by [Amarante \(2019\)](#) can be directly implemented to retrieve nonlinear versions of the Sandwich Theorem, restricting (super/sub)linearity only to some portions of the domain of the functionals involved. Specifically, we define a binary relation (denoted by \mathbf{C}) that we call the *convex-conic symmetric preorder*, which is a symmetric

2020 *Mathematics Subject Classification.* Primary: 46A22; Secondary: 06A06, 06F20.

Key words and phrases. Sandwich Theorem; Pataia's Theorem; \mathbf{C} -(super/sub)linearity; convex-conic symmetric preorder.

We thank Andrea Aveni, Fabio Maccheroni, and René Pfitscher for useful discussions, as well as Stephen Simons for providing some relevant references. Mario Ghossoub acknowledges financial support from the Natural Sciences and Engineering Research Council of Canada (NSERC) (Grant No. 2018-03961). Giulio Principi is grateful for the financial support provided by the Henry M. MacCracken Fellowship at New York University.

preorder closed with respect to positive scalar multiplication and addition. Using this preorder, we provide a Sandwich Theorem (see Theorem 2) for \mathbf{C} -(super/sub)linear functionals, i.e., functionals that satisfy positive homogeneity and (super/sub)additivity only with respect to $(x, y) \in \mathbf{C}$. We then use this result to prove a Hahn-Banach type extension result (see Corollary 1) and an envelope representation result (see Corollary 2). We conclude by providing an illustration of the applicability of our results, with a focus on comonotonic subadditive functionals.

2. PATARAIA'S THEOREM AND ORDER-THEORETIC TERMINOLOGY

2.1. Order-Theoretic Terminology. ¹ Fix a nonempty set S . By a *(binary) relation* over S we mean a set $\mathbf{R} \subseteq S \times S$. For all $x, y, z \in S$, we will often write $x\mathbf{R}y$ in place of $(x, y) \in \mathbf{R}$, and $x\mathbf{R}y\mathbf{R}z$ instead of $x\mathbf{R}y$ and $y\mathbf{R}z$.

We say that a binary relation \mathbf{R} is *reflexive* if $x\mathbf{R}x$ for all $x \in S$, while it is *symmetric* if $x\mathbf{R}y$ implies $y\mathbf{R}x$, for all $x, y \in S$. If $x\mathbf{R}y$ and $y\mathbf{R}z$ implies $x\mathbf{R}z$, for all $x, y, z \in S$, then \mathbf{R} is said to be *transitive*. A reflexive and transitive relation is called a *preorder*. A preorder \mathbf{R} is a *partial order* if it is also *antisymmetric*, that is, if $x\mathbf{R}y$ and $y\mathbf{R}x$, then $x = y$, for all $x, y \in S$, in which case we say that (S, \mathbf{R}) is a *partially ordered set* (henceforth, *poset*). A partial order \mathbf{R} is said to be a *total order* if for all $x, y \in S$, either $x\mathbf{R}y$ or $y\mathbf{R}x$, in which case we say that (S, \mathbf{R}) is a *totally ordered set*. A totally ordered subset of a poset will be referred to as a *chain*.

Given a poset (S, \mathbf{R}) we say that an element $x \in S$ is an \mathbf{R} -upper bound of $B \subseteq S$ if $x\mathbf{R}B$, that is, $x\mathbf{R}y$ for all $y \in B$. We say that x is an \mathbf{R} -supremum for $B \subseteq S$ if it is an \mathbf{R} -upper bound of B and $z\mathbf{R}x$, for all $z \in S$ with $z\mathbf{R}B$. Infima are defined analogously. Given a poset (S, \mathbf{R}) , we denote \mathbf{R} -suprema and \mathbf{R} -infima of $B \subseteq S$ by $\mathbf{R}\text{-sup } B$ and $\mathbf{R}\text{-inf } B$, respectively. When the binary relation is well-understood within the context we will simply write \sup and \inf to ease the notation. We say that a poset (S, \mathbf{R}) is *strictly inductively ordered* if every chain $C \subseteq S$ admits a \mathbf{R} -supremum. Given a poset (S, \mathbf{R}) , we say that a mapping $F : S \rightarrow S$ is \mathbf{R} -inflationary if $F(s)\mathbf{R}s$, for all $s \in S$. Given a nonempty set A , a poset (S, \mathbf{R}) , and two maps $f, g : A \rightarrow S$, we say that $f\mathbf{R}g$ if and only if $f(a)\mathbf{R}g(a)$ for all $a \in A$. We will refer to this order as the \mathbf{R} -pointwise order, and, often simply as the *pointwise order*, when the underlying relation is clear from the context. It is immediately noticeable that given a poset (S, \mathbf{R}) and a set A , the induced \mathbf{R} -pointwise order on S^A is also a partial order.

We say that a (real) vector space V is an *ordered vector space* if it is endowed with a partial order \leq , such that $x \leq y$ if and only if $x + z \leq y + z$ and $\alpha x \leq \alpha y$, for all $x, y, z \in V$ and all $\alpha \geq 0$.² In addition, we say that an ordered vector space (V, \leq) is a *Riesz space* if it is also a *lattice*, that is, $\sup \{x, y\}, \inf \{x, y\} \in V$, for all $x, y \in V$. A subset B of a Riesz space (V, \leq) is said to be *bounded from above* if it admits an upper bound in V . Finally, we say that

¹We summarize in this subsection all the order-theoretic concepts that we use in this paper. For a comprehensive treatment of these notions, we refer the reader to Caspard et al. (2012) and Schröder (2016).

²We will exclusively focus on real vector spaces, which we will henceforth refer to simply as vector spaces.

a Riesz space (V, \geq) is *Dedekind-complete* if each subset of V which is bounded from above admits a supremum.³ For any vector space V we will denote by $\mathbf{0}_V$ its null element.

2.2. Patarai's Fixed Point Theorem. Similarly to [Amarante \(2019\)](#) we will apply the following version of Patarai's Fixed Point Theorem (see also [Escardó \(2003\)](#)).

Theorem 1 (Patarai). *Let (S, \mathbf{R}) be a poset and \mathcal{I} be the set of all inflationary mappings on S . If (S, \mathbf{R}) is strictly inductively ordered, then \mathcal{I} has a common fixed point, that is, there exists $x \in S$ such that $f(x) = x$, for every $f \in \mathcal{I}$.*

Proof. Since \mathbf{R} is reflexive, $\text{Id} \in \mathcal{I}$. Moreover, given that all $f \in \mathcal{I}$ are inflationary, it follows that $f\mathbf{R}\text{Id}$. Suppose that \mathcal{C} is a chain in \mathcal{I} and define $\bar{f} : s \mapsto \sup \{f(s) : f \in \mathcal{C}\}$. Clearly $\bar{f} \in \mathcal{I}$ and hence it is a \mathbf{R} -supremum for \mathcal{C} with respect to the pointwise order. This yields that \mathcal{I} with the pointwise order is strictly inductively ordered. Therefore, by Zorn's Lemma, \mathcal{I} has a maximal element, say $M \in \mathcal{I}$. Note that for all $s \in S$ and all $f \in \mathcal{I}$, $f(M(s))\mathbf{R}M(s)$. Thus, since M is maximal and \mathbf{R} is antisymmetric, we must have $f \circ M = M$. Therefore, for all $s \in S$, $M(s)$ is a common fixed point of \mathcal{I} . \square

3. MAIN RESULTS

3.1. Convex-Conic Symmetric Preorders. Given a vector space V , we say that $X \subseteq V$ is a *convex cone* if $\lambda X \subseteq X$ for all $\lambda > 0$, and $X + X \subseteq X$. Now let X be a given convex cone.

Definition 1. A binary relation $\mathbf{C} \subseteq X \times X$ is a *convex-conic symmetric preorder* if it is reflexive, transitive, symmetric, and it satisfies the following properties:

- (i) $(\lambda x)\mathbf{C}x$, for all $x \in X$ and $\lambda > 0$.
- (ii) $x\mathbf{C}y\mathbf{C}z$ implies $(x + y)\mathbf{C}z$, for all $x, y, z \in X$.

The “convex-conic” adjective in the definition of a convex-conic symmetric preorder is due to the fact that $\mathbf{C}(x) = \{y \in X : y\mathbf{C}x\}$ is a convex cone. Note that the symmetry property (ii) in Definition 1 implies that whenever $x\mathbf{C}y\mathbf{C}z$, we have $(g + u)\mathbf{C}h$, for all $g, u, h \in \{x, y, z\}$. It is important to observe that whenever $\{\mathbf{0}_V\} \times X \subseteq \mathbf{C}$, for some convex-conic symmetric preorder \mathbf{C} on X , it follows that $X \times X = \mathbf{C}$. This is a straightforward consequence of symmetry and transitivity. Because of property (ii), the same would hold if $(-x)\mathbf{C}x$ for all $x \in X$ with $-x \in X$. For future reference we recollect these simple observations on the following lemma.

Lemma 1. *Let X be a convex cone and $\mathbf{C} \subseteq X \times X$ a convex-conic symmetric preorder. If either $\{\mathbf{0}_V\} \times X \subseteq \mathbf{C}$ or $(-x)\mathbf{C}x$ for all $x \in X$ with $-x \in X$, then $X \times X = \mathbf{C}$.*

Therefore, at first sight one may worry about how permissive a convex-conic symmetric preorder is. We show by means of examples (see Section 3.2.2) that it is neither always trivial nor too restrictive.

³We refer to [Aliprantis and Burkinshaw \(2006\)](#) for a detailed treatment of Riesz spaces.

3.2. A Sandwich Theorem. Let (\mathbb{V}, \leq) be a Dedekind-complete Riesz space. Following [Fuchssteiner and Wright \(1977\)](#), we adjoin to \mathbb{V} an element denoted by $-\infty$, and we extend \leq to $\mathbb{V} \cup \{-\infty\}$ assuming $-\infty \leq \mathbb{V}$. We say that a map $F : X \rightarrow \mathbb{V} \cup \{-\infty\}$ is **C-sublinear** if

- (i) F is *positively homogeneous*, i.e., $F(\lambda x) = \lambda F(x)$, for all $\lambda > 0$ and all $x \in X$.
- (ii) F is **C-subadditive**, i.e., $F(x + y) \leq F(x) + F(y)$, for all $x, y \in X$ with $x \mathbf{C} y$.

The definitions of **C-superlinearity** and **C-linearity** are analogous. We endow V with a partial order \preceq such that (V, \preceq) is a partially ordered vector space. We say that a function $F : X \rightarrow \mathbb{V} \cup \{-\infty\}$ is *monotone* if $x \preceq y$ implies $F(x) \leq F(y)$, for all $x, y \in X$. We adopt the convention that $0 \cdot (-\infty) = \mathbf{0}_{\mathbb{V}}$. We are now ready to state and prove our main result.

Theorem 2. *Suppose that $P : X \rightarrow \mathbb{V} \cup \{-\infty\}$ is C-superlinear and $H : X \rightarrow \mathbb{V} \cup \{-\infty\}$ is C-sublinear and monotone. If $P \leq H$, then there exists a C-linear map $Q : X \rightarrow \mathbb{V} \cup \{-\infty\}$ such that $P \leq Q \leq H$.*

The proof will be provided in several steps. Before going into its details, we provide some definitions and simple remarks. Let

$$\mathcal{D}_{\text{PH}} = \{Q : X \rightarrow \mathbb{V} \cup \{-\infty\} : Q \text{ is C-superlinear and } P \leq Q \leq H\}.$$

Clearly, $P \in \mathcal{D}_{\text{PH}} \neq \emptyset$. Moreover, \mathcal{D}_{PH} is a poset with respect to the pointwise order. If \mathcal{C} is a chain in \mathcal{D}_{PH} , then $\overline{Q} : x \mapsto \sup \{Q(x) : Q \in \mathcal{C}\}$ is C-superlinear and a supremum of \mathcal{C} . Indeed, for all $(x, y) \in \mathbf{C}$ and all $Q \in \mathcal{C}$,

$$\overline{Q}(x + y) \geq Q(x + y) \geq Q(x) + Q(y).$$

This implies that \overline{Q} is C-superlinear. Thus \mathcal{D}_{PH} is strictly inductively ordered. Let $A : Q \mapsto A_Q$ be defined as

$$A_Q(x) = \inf_y \{H(x + y) - Q(y) : y \in \mathbf{C}(x) \text{ and } Q(y) > -\infty\},$$

for all $x \in X$ and all $Q \in \mathcal{D}_{\text{PH}}$. Now fix $g, x \in X$ and $Q \in \mathcal{D}_{\text{PH}}$. We also define,

$$T_g(Q)(x) = \sup_{h, \lambda} \{Q(h) + \lambda A_Q(g) : h + \lambda g \preceq x, h \in \mathbf{C}(x), \lambda \geq 0\}.$$

We then obtain the following result.

Lemma 2. *The following claims hold:*

- (1) For all $Q \in \mathcal{D}_{\text{PH}}$, we have $A_Q \geq Q$ and A_Q is C-sublinear.
- (2) For all $Q \in \mathcal{D}_{\text{PH}}$ and $g \in X$, we have $T_g(Q)$ is C-superlinear.
- (3) For all $Q \in \mathcal{D}_{\text{PH}}$, we have $Q \leq T_g(Q) \leq H$.
- (4) For all $Q \in \mathcal{D}_{\text{PH}}$ and $x \in X$, we have $T_x(Q)(x) \geq A_Q(x)$.

Proof. (1). Fix $Q \in \mathcal{D}_{\text{PH}}$ and $x \in X$, then

$$H(x + y) - Q(y) \geq Q(x + y) - Q(y) \geq Q(x) + Q(y) - Q(y) = Q(x),$$

for all $y \in \mathbf{C}(x)$ with $Q(y) > -\infty$. Therefore, $A_Q \geq Q$. Now suppose that $(x_1, x_2) \in \mathbf{C}$ and $y_1, y_2 \in \mathbf{C}(x_1)$ with $Q(y_1) > -\infty$, $Q(y_2) > -\infty$. Notice that since $y_1, y_2 \in \mathbf{C}(x_1)$ and \mathbf{C} is symmetric and transitive, we have $x_1 \mathbf{C} y_1 \mathbf{C} y_2 \mathbf{C} x_2$. By symmetry and property (ii), it follows that

$$((x_1 + y_1), (x_2 + y_2)) \in \mathbf{C}.$$

These observations imply that $Q(y_1 + y_2) \geq Q(y_1) + Q(y_2)$, and hence

$$\begin{aligned} H(x_1 + x_2 + y_1 + y_2) - Q(y_1 + y_2) &\leq H(x_1 + x_2 + y_1 + y_2) - Q(y_1) - Q(y_2) \\ &\leq H(x_1 + y_1) - Q(y_1) + H(x_2 + y_2) - Q(y_2). \end{aligned}$$

Therefore, $A_Q(x_1 + x_2) \leq A_Q(x_1) + A_Q(x_2)$. Consequently, for all $Q \in \mathcal{D}_{\text{PH}}$, A_Q is \mathbf{C} -subadditive. Positive homogeneity follows from the positive homogeneity of Q and H , as well as from the fact that $\mathbf{C}(x)$ is a cone.

(2). Fix $Q \in \mathcal{D}_{\text{PH}}$, $g \in X$, and $(x_1, x_2) \in \mathbf{C}$. Then $T_g(Q)(x_1 + x_2) \geq T_g(Q)(x_1) + T_g(Q)(x_2)$. Indeed, notice that if $h_1 \in \mathbf{C}(x_1)$, $h_2 \in \mathbf{C}(x_2)$, $\lambda_1, \lambda_2 \geq 0$ satisfy

$$h_1 + \lambda_1 g \preceq x_1 \text{ and } h_2 + \lambda_2 g \preceq x_2,$$

then $h_1 + h_2 + (\lambda_1 + \lambda_2)g \preceq x_1 + x_2$. Since $x_1 \mathbf{C} x_2$, by symmetry, transitivity, and property (ii) we have that $x_1 \mathbf{C} x_2 \mathbf{C} h_1 \mathbf{C} h_2$ and

$$(h_1 + h_2, \lambda_1 + \lambda_2) \in \{(h, \lambda) : h + \lambda g \preceq x_1 + x_2, h \in \mathbf{C}(x_1 + x_2), \lambda \geq 0\}.$$

This, together with the \mathbf{C} -superadditivity of Q , yields

$$\begin{aligned} T_g(Q)(x_1 + x_2) &= \sup_{h, \lambda} \{Q(h) + \lambda A_Q(g) : h + \lambda g \preceq x_1 + x_2, h \in \mathbf{C}(x_1 + x_2), \lambda \geq 0\} \\ &\geq \sup_{h_1, h_2, \lambda_1, \lambda_2} \left\{ Q(h_1 + h_2) + (\lambda_1 + \lambda_2) A_Q(g) : \begin{array}{l} h_1 + h_2 + (\lambda_1 + \lambda_2)g \preceq x_1 + x_2, \\ h_1 \in \mathbf{C}(x_1), h_2 \in \mathbf{C}(x_2), \lambda_1, \lambda_2 \geq 0 \end{array} \right\} \\ &\geq Q(h_1) + Q(h_2) + \lambda_1 A_Q(g) + \lambda_2 A_Q(g), \end{aligned}$$

for all $h_1 \in \mathbf{C}(x_1)$, $h_2 \in \mathbf{C}(x_2)$, $\lambda_1, \lambda_2 \geq 0$ with $h_1 + \lambda_1 g \preceq x_1$ and $h_2 + \lambda_2 g \preceq x_2$. Thus,

$$T_g(Q)(x_1 + x_2) \geq T_g(Q)(x_1) + T_g(Q)(x_2).$$

This proves that for all $Q \in \mathcal{D}_{\text{PH}}$ and all $g \in X$, the mapping $T_g(Q)$ is \mathbf{C} -superadditive. Positive homogeneity of each $T_g(Q)$ follows from the positive homogeneity of Q and A_Q (see the previous point) and the fact that $\mathbf{C}(x)$ is a convex cone.

(3). Fix $Q \in \mathcal{D}_{\text{PH}}$. It is immediate to see that $T_g(Q) \geq Q$ (take $\lambda = 0$ and $h = x$, recalling that \mathbf{C} is reflexive). We show that $T_g(Q) \leq H$. First notice that, if $\lambda = 0$, then for all $x \in X$, by the monotonicity of H ,

$$\sup_h \{Q(h) + \lambda A_Q(g) : h + \lambda g \preceq x, h \in \mathbf{C}(x)\} = \sup_h \{Q(h) : h \preceq x, h \in \mathbf{C}(x)\}$$

$$\leq \sup_h \{H(h) : h \preceq x, h \in \mathbf{C}(x)\} \leq H(x).$$

Therefore, we can focus on all $\lambda > 0$. In particular, we have that for all $x \in X$,

$$\begin{aligned} & \sup_{h,\lambda} \{Q(h) + \lambda A_Q(g) : h + \lambda g \preceq x, h \in \mathbf{C}(x), \lambda > 0\} \\ &= \sup_{h,\lambda} \inf_{y \in \mathbf{C}(x), Q(y) > -\infty} \{Q(h) + \lambda H(g+y) - \lambda Q(y) : h + \lambda g \preceq x, h \in \mathbf{C}(x), \lambda > 0\} \\ &= \sup_{h,\lambda} \inf_{y \in \mathbf{C}(x), Q(y) > -\infty} \{Q(h) + H(\lambda g + \lambda y) - Q(\lambda y) : h + \lambda g \preceq x, h \in \mathbf{C}(x), \lambda > 0\} \\ &\leq \sup_{h,\lambda} \{H(\lambda g + h) : h + \lambda g \preceq x, h \in \mathbf{C}(x), \lambda > 0\} \\ &= \sup_{h,\lambda} \{H(\lambda g + \lambda h) : \lambda h + \lambda g \preceq x, h \in \mathbf{C}(x), \lambda > 0\} \\ &\leq H(x), \end{aligned}$$

where the last two steps follow from the fact that $\mathbf{C}(x)$ is a cone and H is monotone. Thus, connecting the two observations for $\lambda = 0$ and all $\lambda > 0$, we have that

$$T_g(Q)(x) = \sup_{h,\lambda} \{Q(h) + \lambda A_Q(g) : h + \lambda g \preceq x, h \in \mathbf{C}(x), \lambda \geq 0\} \leq H(x),$$

for all $x \in X$, and hence $Q \leq T_g(Q) \leq H$.

(4). Let $x \in X$ and $Q \in \mathcal{D}_{\text{PH}}$. Then, since Q is positively homogeneous, we have

$$\begin{aligned} T_x(Q)(x) &= \sup_{h,\lambda} \{Q(h) + \lambda A_Q(x) : h + \lambda x \preceq x, h \in \mathbf{C}(x), \lambda \geq 0\} \\ &\geq \sup_{h,n} \left\{ Q(h) + \frac{n-1}{n} A_Q(x) : h \preceq \frac{1}{n}x, h \in \mathbf{C}(x) \right\} \\ &\geq \frac{1}{m} Q(x) + \frac{m-1}{m} A_Q(x) \\ &\geq -\frac{1}{m} |Q(x)| + \frac{m-1}{m} A_Q(x), \end{aligned}$$

for all $m \in \mathbb{N}$. Thus, letting $m \rightarrow \infty$, since all Dedekind complete Riesz spaces are Archimedean⁴ (see Lemma 8.4 in Aliprantis and Border (2006)), we have $T_x(Q)(x) \geq A_Q(x)$ for all $x \in X$. \square

This lemma highlights the fact that for all $g \in X$, $T_g(\cdot)$ is a selfmap, as $T_g(Q) \in \mathcal{D}_{\text{PH}}$ for all $Q \in \mathcal{D}_{\text{PH}}$. Now we are ready to provide the proof of Theorem 2.

Proof of Theorem 2. By Lemma 2, for all $g \in X$, $T_g : \mathcal{D}_{\text{PH}} \rightarrow \mathcal{D}_{\text{PH}}$ is inflationary. Therefore $(T_g)_{g \in X}$ is a family of inflationary functions. By Theorem 1 the family $(T_g)_{g \in X}$ has a common fixed point $Q^* \in \mathcal{D}_{\text{PH}}$. Since for all $g \in X$ and all $Q \in \mathcal{D}_{\text{PH}}$ we have $T_g(Q) \leq H$, it follows that for all $x \in X$ and $y \in \mathbf{C}(x)$ with $Q^*(y) > -\infty$,

$$\begin{aligned} H(x+y) - Q^*(y) &\geq T_g(Q^*)(x+y) - Q^*(y) \\ &\geq T_g(Q^*)(x) + T_g(Q^*)(y) - T_g(Q^*)(y) = T_g(Q^*)(x). \end{aligned}$$

⁴We recall that a Riesz space (W, \leq) is *Archimedean* if whenever $\mathbf{0}_W \leq nx \leq y$ for all $n = 1, 2, \dots$ and some $y \geq \mathbf{0}_W$, we have that $x = \mathbf{0}_W$.

Thus, $A_{Q^*}(x) \geq T_g(Q^*)(x) = Q^*(x)$ for all $x, g \in X$. Moreover, by Lemma 2-(4), we have $T_g(Q^*)(x) = Q^*(x) = T_x(Q^*)(x) \geq A_{Q^*}(x)$, for all $x, g \in X$. Therefore, $A_{Q^*} = Q^*$. Since A_{Q^*} is \mathbf{C} -sublinear and Q^* is \mathbf{C} -superlinear we have that Q^* is \mathbf{C} -linear. To conclude, since $Q^* \in \mathcal{D}_{\text{PH}}$, the claim follows. \square

3.2.1. Extension and Envelope Results. By Theorem 2, we derive an analogous version of the Hahn-Banach Extension Theorem (see for example Theorem 1.25 in Aliprantis and Burkinshaw (2006)).

Corollary 1. *Let $H : X \rightarrow \mathbb{V} \cup \{-\infty\}$ be \mathbf{C} -sublinear and monotone, and $Y \subseteq X$ be a convex cone. If $\ell : Y \rightarrow \mathbb{V} \cup \{-\infty\}$ is \mathbf{C} -linear and satisfies $\ell \leq H|_Y$, then there exists a \mathbf{C} -linear map $Q : X \rightarrow \mathbb{V} \cup \{-\infty\}$ such that $Q \leq H$ and $\ell \leq Q|_Y$.*

Proof. Define $P : X \rightarrow \mathbb{V} \cup \{-\infty\}$ by

$$P(x) = \begin{cases} \ell(x) & x \in Y, \\ -\infty & x \notin Y, \end{cases}$$

for all $x \in X$. Thus P is \mathbf{C} -superadditive. Indeed, suppose that $x \mathbf{C} y$. If $x, y \in Y$, then there is nothing to prove. If $x \notin Y$ or $y \notin Y$, then $P(x) + P(y) = -\infty \leq P(x + y)$. It is immediate to see that P is positively homogeneous and $P \leq H$. Therefore, by Theorem 2, there exists a \mathbf{C} -linear map $Q : X \rightarrow \mathbb{V} \cup \{-\infty\}$ such that $P \leq Q \leq H$. Therefore, $\ell = P|_Y \leq Q|_Y$. \square

Next, as a direct application of Corollary 1, we provide a general envelope representation result for \mathbf{C} -sublinear and monotone maps. For all $x \in X$, denote by C_x the convex cone generated by x , that is, $C_x = \{\lambda x : \lambda > 0\}$. Moreover, for all maps $H : X \rightarrow \mathbb{V} \cup \{-\infty\}$, let

$$D(H) = \{Q : X \rightarrow \mathbb{V} \cup \{-\infty\} : Q \text{ is } \mathbf{C}\text{-linear and } Q \leq H\}.$$

Corollary 2. *If $H : X \rightarrow \mathbb{V} \cup \{-\infty\}$ is \mathbf{C} -sublinear and monotone, then*

$$H(x) = \sup_{Q \in D(H)} Q(x), \text{ for all } x \in X.$$

Proof. Let $x \in X$ and define ℓ over C_x as $\ell(\lambda x) = \lambda H(x)$, for all $\lambda x \in C_x$ with $\lambda > 0$. Notice that ℓ is \mathbf{C} -linear and positively homogeneous. Moreover, by positive homogeneity we have that $\ell(\lambda x) = \lambda H(x) = H(\lambda x)$ for all $\lambda x \in C_x$. Thus, by Corollary 1, there exists a \mathbf{C} -linear map $Q : X \rightarrow \mathbb{V} \cup \{-\infty\}$ such that $Q \leq H$ and $\ell \leq Q|_{C_x}$. Moreover,

$$H(x) = \ell(x) \leq Q(x) \leq H(x),$$

and so $Q(x) = H(x)$. Therefore, for all $x \in X$, there exists a \mathbf{C} -linear map $Q_x : X \rightarrow \mathbb{V} \cup \{-\infty\}$ such that $Q_x(x) = H(x)$. This implies that

$$H(x) = \sup_{Q \in D(H)} Q(x),$$

for all $x \in X$. \square

Remark 1. Suppose that $\mathbb{V} = \mathbb{R}$ and V is a topological vector space. For any map $G : V \rightarrow [-\infty, \infty)$, we define the (effective) domain of G by $\text{dom}G = \{x \in V : G(x) > -\infty\}$. We say that G is proper if $\text{dom}G$ is nonempty. It is important to note that Corollary 2 could also be proved by defining each ℓ on the vector space generated by $x \in V$, i.e., $\text{span}\{x\}$, as $\ell(\alpha x) = \alpha H(x)$ for all $\alpha \in \mathbb{R}$, and then extending each ℓ to be equal to $-\infty$ everywhere else. In such a case, we would have that each extension $Q|_{\text{dom}G}$ is continuous. Therefore, in this setting, Corollary 2 could be restated, adding a further restriction on $D(H)$, namely the continuity of its elements when restricted to their domains. This approach is similar to the one adopted by Roth (2000) (e.g., see Corollary 3.3).

3.2.2. Examples.

Example 1. Clearly $\mathbf{C} = X \times X$ is a convex-conic symmetric preorder. Under this convex-conic symmetric preorder \mathbf{C} -(super/sub)linearity corresponds to the classic (super/sub)linearity.

Example 2. [Positive homogeneity] Fix $x \in X$ and define $\mathbf{D}_x = \{(\lambda x, \beta x) : \lambda, \beta > 0\}$. Then \mathbf{D}_x is a convex-conic symmetric preorder. More importantly, define $\mathbf{D} = \bigcup_{x \in X} \mathbf{D}_x$. We now verify that \mathbf{D} is also a convex-conic symmetric preorder. Since each $\mathbf{D}_x \subseteq \mathbf{D}$, we have that \mathbf{D} is symmetric, reflexive, and satisfies property (i). Fix arbitrarily $x, y, z \in X$ and suppose that $x \mathbf{D} y \mathbf{D} z$. Then there exist $\alpha, \beta, \lambda, \gamma > 0$ and $v, w \in X$ such that,

$$(x, y) = (\alpha v, \beta v) \text{ and } (y, z) = (\lambda w, \gamma w).$$

Therefore, $x = \alpha v = \frac{\alpha}{\beta} y = \lambda \frac{\alpha}{\beta} w$, and hence $x \mathbf{D} z$. Consequently, \mathbf{D} is transitive. Moreover,

$$(x + y, z) = \left(\lambda \left(\frac{\alpha}{\beta} + 1 \right) w, \gamma w \right),$$

and thus \mathbf{D} satisfies property (ii). Note that any positively homogeneous map $F : X \rightarrow \mathbb{R}$ is \mathbf{D} -linear.

Example 3. [Equivalent measures] Consider a measurable space (Ω, \mathcal{F}) and denote by X the convex cone of countably additive measures $\mu : \mathcal{F} \rightarrow [0, \infty]$. For all measures $\mu \in X$ we define $N_\mu = \{A \in \mathcal{F} : \mu(A) = 0\}$, i.e., the collection of μ -null elements of \mathcal{F} . Two measures $\mu, \nu \in X$ are said to be *equivalent*, denoted by $\mu \sim \nu$, if $N_\mu = N_\nu$. It can be verified that \sim is a convex-conic symmetric preorder. Indeed, letting $\lambda > 0$ and $\mu \in X$, we have $N_{\lambda\mu} = N_\mu$. If $\nu, \mu, \eta \in X$, $\nu \sim \mu$, and $\mu \sim \eta$, then $N_\mu = N_\nu = N_\eta$. Thus \sim is transitive. Symmetry is also straightforward. To conclude, if $\nu \sim \mu \sim \eta$ for $\nu, \mu, \eta \in X$, we have that $\nu(A) + \mu(A) = 0$ for all $A \in N_\eta$, and the converse holds as well. This yields $N_{\nu+\mu} = N_\eta$, and hence \sim is a convex-conic symmetric preorder.

Example 4. [Strict comonotonicity] Let (Ω, \mathcal{F}) be a measurable space, and denote by $L^0(\Omega, \mathcal{F})$ the space of \mathcal{F} -measurable real-valued functions. For all $x \in L^0(\Omega, \mathcal{F})$, denote by C_x the convex cone generated by x . We say that $x, y \in L^0(\Omega, \mathcal{F})$ are *strictly comonotonic* if either (1) $x \in C_y$; or (2) for all $\omega_1 \neq \omega_2$ in Ω ,

$$[x(\omega_1) - x(\omega_2)] [y(\omega_1) - y(\omega_2)] > 0.$$

The strict comonotonicity relation, denoted as $\uparrow\uparrow$, requires that $x \uparrow\uparrow y$ if and only if $x, y \in L^0(\Omega, \mathcal{F})$ are strictly comonotonic. We now verify that $\uparrow\uparrow$ is a convex-conic symmetric order. Clearly, $(\lambda x) \uparrow\uparrow x$, for all $\lambda > 0$, as $\lambda x \in C_x$, for all $x \in L^0(\Omega, \mathcal{F})$. Symmetry is immediate, since if $x \in C_y$, then $y \in C_x$. To show transitivity, fix $\omega_1 \neq \omega_2$, and suppose that $x \uparrow\uparrow y$ and $y \uparrow\uparrow z$. Then, we have three cases:

(1) If $x \in C_y$ and $y \in C_z$, then $x \in C_z$. Thus, $x \uparrow\uparrow z$.

(2) If $x \in C_y$ and $y \notin C_z$, then

$$[y(\omega_1) - y(\omega_2)] [z(\omega_1) - z(\omega_2)] > 0 \quad \text{and} \quad x = \lambda y,$$

for some $\lambda > 0$. This implies that

$$[x(\omega_1) - x(\omega_2)] [z(\omega_1) - z(\omega_2)] > 0.$$

Thus, $x \uparrow\uparrow z$.

(3) If $x \notin C_y$ and $y \notin C_z$, then

$$[x(\omega_1) - x(\omega_2)] [y(\omega_1) - y(\omega_2)] > 0 \quad \text{and} \quad [y(\omega_1) - y(\omega_2)] [z(\omega_1) - z(\omega_2)] > 0.$$

Therefore, if $x(\omega_1) - x(\omega_2) > 0$, then $y(\omega_1) - y(\omega_2) > 0$, and hence $z(\omega_1) - z(\omega_2) > 0$.

The same reasoning (with inverted signs) applies to the case where $x(\omega_1) - x(\omega_2) < 0$.

Thus $x \uparrow\uparrow z$.

To conclude, fix $\omega_1 \neq \omega_2$ and suppose that $x \uparrow\uparrow y \uparrow\uparrow z$. Then, we have three cases:

(1) If $x \in C_y$ and $y \in C_z$, then $x + y \in C_z$. Thus, $(x + y) \uparrow\uparrow z$.

(2) If $x \in C_y$ and $y \notin C_z$, then

$$[y(\omega_1) - y(\omega_2)] [z(\omega_1) - z(\omega_2)] > 0 \quad \text{and} \quad x = \lambda y,$$

for some $\lambda > 0$. This implies that

$$[x(\omega_1) + y(\omega_1) - x(\omega_2) - y(\omega_2)] [z(\omega_1) - z(\omega_2)] > 0.$$

Thus, $(x + y) \uparrow\uparrow z$.

(3) If $x \notin C_y$ and $y \notin C_z$, then

$$\begin{aligned} & [x(\omega_1) + y(\omega_1) - x(\omega_2) - y(\omega_2)] [z(\omega_1) - z(\omega_2)] \\ &= [x(\omega_1) - x(\omega_2)] [z(\omega_1) - z(\omega_2)] + [y(\omega_1) - y(\omega_2)] [z(\omega_1) - z(\omega_2)] > 0. \end{aligned}$$

Thus, $(x + y) \uparrow\uparrow z$.

Therefore, $\uparrow\uparrow$ is a convex-conic symmetric preorder.

Comonotonicity is an important property in decision theory, risk measurement, and the theory of risk sharing. A different example of a convex-conic preorder concerns vectors that are affinely related.

Example 5. [Affinity] Let X be a vector space, and fix $e \in X$. We say that x and y in X are *e-affinely related* if there exist $\alpha, \beta \neq 0$ such that $x = \alpha y + \beta e$. In particular, we define

the binary relation **Aff** as follows

$$x\mathbf{Aff}y \iff \exists \alpha \neq 0, \exists \beta \in \mathbb{R} : x = \alpha y + \beta e.$$

Even if straightforward, we provide the steps proving that **Aff** is a convex-conic symmetric preorder. Reflexivity is immediate, as $x = x$. Passing to symmetry, if $x\mathbf{Aff}y$, then there exist $\alpha \neq 0$ and $\beta \in \mathbb{R}$ such that $x = \alpha y + \beta e$, and hence $y = \frac{1}{\alpha}x - \beta e$, implying that $y\mathbf{Aff}x$. Suppose that $x\mathbf{Aff}y$ and $y\mathbf{Aff}z$. There exist $\alpha_1, \alpha_2 \neq 0$ and $\beta_1, \beta_2 \in \mathbb{R}$ such that $x = \alpha_1 y + \beta_1 e$ and $y = \alpha_2 z + \beta_2 e$. Then

$$x = \alpha_1(\alpha_2 z + \beta_2 e) + \beta_1 e = \alpha_1 \alpha_2 z + (\alpha_1 \beta_2 + \beta_1) e,$$

proving that $x\mathbf{Aff}z$, and so **Aff** is transitive. Now fix $x \in X$, then clearly $(\lambda x)\mathbf{Aff}x$ for all $\lambda > 0$. In conclusion, if $x\mathbf{Aff}y\mathbf{Aff}z$, then there exist $\alpha_1, \alpha_2 \neq 0$ and $\beta_1, \beta_2 \in \mathbb{R}$ such that $x = \alpha_1 y + \beta_1 e$ and $y = \alpha_2 z + \beta_2 e$. Then

$$\begin{aligned} x + y &= \alpha_1 y + \beta_1 e + \alpha_2 z + \beta_2 e = \alpha_1 \alpha_2 z + \alpha_1 \beta_2 e + \beta_1 e + \alpha_2 z + \beta_2 e \\ &= (\alpha_1 \alpha_2 + \alpha_2) z + (\alpha_1 \beta_2 + \beta_1 + \beta_2) e, \end{aligned}$$

implying that $(x + y)\mathbf{Aff}z$. Thus **Aff** is a convex-conic symmetric preorder.

To conclude this section we provide some examples of binary relations that are *not* convex-conic symmetric preorders. Clearly any irreflexive, asymmetric, or nontransitive binary relation would work. For instance, probabilistic independence and orthogonality in the context of inner product spaces are relevant examples.

Example 6. [Inner product spaces] Consider the Hilbert space $L^2([0, 1], \mathcal{B}[0, 1], \text{Leb})$, where $\mathcal{B}[0, 1]$ denotes the Borel sigma algebra and Leb the Lebesgue measure. We define the relation **Corr** by

$$f\mathbf{Corr}g \iff \int_{[0,1]} fg \, d\text{Leb} \geq 0.$$

It is easy to see that **Corr** is reflexive and symmetric, and it satisfies conditions (i) and (ii) of Definition 1, but it fails transitivity. Indeed, consider the functions

$$f = \mathbf{1}_{[0,1]}, \quad g = \mathbf{1}_{[0, \frac{1}{2}]}, \quad h = -\mathbf{1}_{[\frac{1}{2}, 1]}.$$

Then,

$$f\mathbf{Corr}g \text{ and } g\mathbf{Corr}h, \text{ but } \int_{[0,1]} fh \, d\text{Leb} = -\frac{1}{2} < 0.$$

The next example provides a symmetric preorder that fails property (ii) of Definition 1.

Example 7. Define the function $\varphi : \mathbb{R}^2 \rightarrow \mathbb{R}$ by

$$\varphi(x, y) = \begin{cases} -1 & (x, y) \in ((\mathbb{R} \setminus \{0\}) \times \{0\}) \cup (\{0\} \times (\mathbb{R} \setminus \{0\})) \\ 0 & \text{otherwise.} \end{cases}$$

Consider the binary relation $x\boldsymbol{\varphi}y$ if and only if $\varphi(x, y) \geq 0$. By definition, $\varphi(x, x) = 0$ for all $x \in \mathbb{R}$, and so $\boldsymbol{\varphi}$ is reflexive. Moreover, $\varphi(x, y) = \varphi(y, x)$ for all $x, y \in \mathbb{R}$, proving that $\boldsymbol{\varphi}$ is

symmetric. Additionally, if $x, y, z \in \mathbb{R}$ with $\varphi(x, y) \geq 0$ and $\varphi(y, z) \geq 0$, then it must be the case that either $x, y, z = 0$ or $x, y, z \neq 0$, and hence, $\varphi(x, z) = 0$. Therefore φ is transitive. However, letting $x, z = 1, y = -1$, we obtain

$$\varphi(x, y) \geq 0, \varphi(y, z) \geq 0, \text{ and } \varphi(x + y, z) = \varphi(0, z) < 0.$$

Thus φ is a symmetric preorder that is not a convex-conic symmetric preorder.

4. RELAXING POSITIVE HOMOGENEITY

In this section, we weaken the definition of a convex-conic symmetric preorder by removing property (i) (i.e., positive homogeneity). In particular, we define a new preorder called “summand symmetric preorder”. Fix a vector space V and a subset $X \subseteq V$ such that $X + X \subseteq X$.

Definition 2. A binary relation $\mathbf{S} \subseteq X \times X$ is a *summand symmetric preorder* if it is reflexive, transitive, and symmetric, and it satisfies the following property:

$$x\mathbf{S}y\mathbf{S}z \text{ implies } (x + y)\mathbf{S}z, \text{ for all } x, y, z \in X. \quad (1)$$

Note that in this case $\mathbf{S}(x) = \{y \in X : y\mathbf{S}x\}$ is closed with respect to the summation of its elements. Clearly, if X is a convex cone and \mathbf{C} is a convex-conic symmetric preorder on X , then \mathbf{C} is also a summand symmetric preorder.

Let (\mathbb{V}, \leq) be a Dedekind-complete Riesz space. We say that a map $F : X \rightarrow \mathbb{V} \cup \{-\infty\}$ is *\mathbf{S} -sublinear* if

- (i) F is *positively integer homogeneous*, i.e., $F(nx) = nF(x)$, for all $n \in \mathbb{N}$ and all $x \in X$.
- (ii) F is *\mathbf{S} -subadditive*, i.e., $F(x + y) \leq F(x) + F(y)$, for all $x, y \in X$ with $x\mathbf{S}y$.

The definitions of \mathbf{S} -superlinearity and \mathbf{S} -linearity are analogous. We endow V with a partial order \preceq such that (V, \preceq) is a partially ordered vector space. We say that a map $F : X \rightarrow \mathbb{V} \cup \{-\infty\}$ is *monotone* if $x \preceq y$ implies $F(x) \leq F(y)$, for all $x, y \in X$. We adopt the convention that $0 \cdot (-\infty) = \mathbf{0}_{\mathbb{V}}$. Using the exact same steps as in the proof of Theorem 2, slightly changing the auxiliary maps, we can retrieve the following version of Theorem 2 for \mathbf{S} -(super/sub)linear functionals.

Theorem 3. Suppose that X is a convex cone, $P : X \rightarrow \mathbb{V} \cup \{-\infty\}$ is \mathbf{S} -superlinear, and $H : X \rightarrow \mathbb{V} \cup \{-\infty\}$ is \mathbf{S} -sublinear and monotone. If $P \leq H$, then there exists an \mathbf{S} -linear map $Q : X \rightarrow \mathbb{V} \cup \{-\infty\}$ such that $P \leq Q \leq H$.

We omit the proof of this result as it is almost identical to that of Theorem 2. We simply provide the form of one of the auxiliary maps we used in the previous proofs. In particular, let

$$\mathcal{D}_{\text{PH}} = \{Q : X \rightarrow \mathbb{V} \cup \{-\infty\} : Q \text{ is } \mathbf{S}\text{-superlinear and } P \leq Q \leq H\}.$$

Now, fix $g, x \in X$, and $Q \in \mathcal{D}_{\text{PH}}$. We also define,

$$T_g(Q)(x) = \sup_{h,n} \{Q(h) + nA_Q(g) : h + ng \preceq x, h \in \mathbf{S}(x), n \in \mathbb{N} \cup \{0\}\}.$$

The auxiliary function A_Q is defined as in Section 3.2, using \mathbf{S} instead of \mathbf{C} .

The analogous of Lemma 2 also holds in this setting. Before providing its proof, we require the following simple result.

Lemma 3. *Suppose that X is a convex cone and \mathbf{S} a summand symmetric preorder on X . Then for all $x, y \in X$,*

$$x\mathbf{S}y \implies \frac{x}{n}\mathbf{S}y \text{ for all } n \in \mathbb{N}.$$

Proof. Fix $x, y \in X$ with $x\mathbf{S}y$, and choose $n \in \mathbb{N}$ arbitrarily. Since \mathbf{S} is reflexive, it follows that $x/n\mathbf{S}x/n$. Hence, by property (1), we obtain

$$x = \left(\sum_{k=1}^n \frac{x}{n} \right) \mathbf{S} \frac{x}{n}.$$

Thus, by transitivity and symmetry of \mathbf{S} , we have $x/n\mathbf{S}y$. □

Lemma 4. *The following claims hold:*

- (1) *For all $Q \in \mathcal{D}_{\text{PH}}$, we have $A_Q \geq Q$ and A_Q is \mathbf{S} -sublinear.*
- (2) *For all $Q \in \mathcal{D}_{\text{PH}}$ and $g \in X$, we have $T_g(Q)$ is \mathbf{S} -superlinear.*
- (3) *For all $Q \in \mathcal{D}_{\text{PH}}$, we have $Q \leq T_g(Q) \leq H$.*
- (4) *For all $Q \in \mathcal{D}_{\text{PH}}$ and all $x \in X$, we have $T_x(Q)(x) \geq A_Q(x)$.*

Proof. We provide a proof only for points (3) and (4). For the remaining points, the proofs are identical to those of Lemma 2.

(3). It is immediate to see that $T_g(Q) \geq Q$, for all $Q \in \mathcal{D}_{\text{PH}}$ (take $n = 0$ and $h = x$). We show that $T_g(Q) \leq H$. In particular, we have that for all $x \in X$,

$$\begin{aligned} T_g(Q)(x) &= \sup_{h,n} \{Q(h) + nA_Q(g) : h + ng \preceq x, h \in \mathbf{S}(x), n \in \mathbb{N} \cup \{0\}\} \\ &= \sup_{h,n} \inf_{y \in \mathbf{S}(x), Q(y) > -\infty} \{Q(h) + nH(g+y) - nQ(y) : h + ng \preceq x, h \in \mathbf{S}(x), n \in \mathbb{N} \cup \{0\}\} \\ &= \sup_{h,n} \inf_{y \in \mathbf{S}(x), Q(y) > -\infty} \{Q(h) + H/ng + ny - Q(ny) : h + ng \preceq x, h \in \mathbf{S}(x), n \in \mathbb{N} \cup \{0\}\} \\ &\leq \sup_{h,n} \{H/ng + h : h + ng \preceq x, h \in \mathbf{S}(x), n \in \mathbb{N} \cup \{0\}\} \\ &= \sup_{h,n} \{H/ng + nh : nh + ng \preceq x, h \in \mathbf{S}(x), n \in \mathbb{N} \cup \{0\}\} \end{aligned}$$

$$\leq H(x),$$

where the fourth inequality follows from setting $y = \frac{h}{n}$ (which belongs to $\mathbf{S}(x)$ by Lemma 3), while the last two steps follow from the fact that $\mathbf{S}(x)$ is closed with respect to addition and H is monotone. Thus, $Q \leq T_g(Q) \leq H$.

(4). Let $x \in X$, $Q \in \mathcal{D}_{\text{PH}}$, and $m \in \mathbb{N}$. Then, since Q and $T_x(Q)$ are integer positively homogeneous, we have

$$\begin{aligned} T_x(Q)(x) &= \frac{T_x(Q)((m+1)x)}{m+1} \\ &= \frac{1}{m+1} \sup_{h,n} \{Q(h) + nA_Q(x) : h + nx \preceq (m+1)x, h \in \mathbf{S}(x), n \in \mathbb{N} \cup \{0\}\} \\ &\geq \frac{1}{m+1} \sup_h \{Q(h) + mA_Q(x) : h \preceq x, h \in \mathbf{S}(x)\} \\ &\geq \frac{1}{m+1} Q(x) + \frac{m}{m+1} A_Q(x) \\ &\geq -\frac{1}{m+1} |Q(x)| + \frac{m}{m+1} A_Q(x). \end{aligned}$$

Thus, letting $m \rightarrow \infty$, since all Dedekind complete Riesz spaces are Archimedean (see Lemma 8.4 in Aliprantis and Border (2006)), we have $T_x(Q)(x) \geq A_Q(x)$, for all $x \in X$. \square

The proof of Theorem 3 is now totally analogous to that of Theorem 2, and it is omitted.

5. APPLICATIONS: COMONOTONIC SUBADDITIVITY

In this section, we apply the previous results to the case of comonotonicity on a specific measurable space. Suppose that $\Omega = [0, 1]$ and $\mathcal{F} = \mathcal{B}[0, 1]$, the Borel sigma algebra on the unit interval. Using our results and observations in Example 4, we obtain the following.

Proposition 1. *Suppose that $H : B(\Omega, \mathcal{F}) \rightarrow [-\infty, \infty)$ is $\|\cdot\|_\infty$ -continuous when restricted to its domain, monotone, positively homogeneous, and strictly comonotonic subadditive. Then*

$$H(x) = \sup_{Q \in D(H)} Q(x), \text{ for all } x \in B(\Omega, \mathcal{F}), \quad (2)$$

where $D(H)$ is a set of maps from $B(\Omega, \mathcal{F})$ to $[-\infty, \infty)$ that are comonotonic additive and $\|\cdot\|_\infty$ -continuous when restricted to their domains. Moreover, H is comonotonic subadditive on its domain.

Proof. By Corollary 2, Remark 1, and Example 4, it follows that (2) holds, where $D(H)$ is a convex set of functionals from $B(\Omega, \mathcal{F})$ to $[-\infty, \infty)$ that are strictly comonotonic additive and $\|\cdot\|_\infty$ -continuous when restricted to their domains. By Lemma 7 in the Appendix, all elements of $D(H)$ are comonotonic additive on their domains. Note that Lemma 7 also implies that H is comonotonic subadditive on its domain. \square

Comonotonic additive functionals play a central role in the theory of decision-making under ambiguity. Their study was pioneered by [Schmeidler \(1989\)](#), who also provided a representation of such functionals in terms of Choquet integrals ([Schmeidler \(1986\)](#)). Such functionals are also relevant for their connection to the theory of risk measurement (e.g., see [Denuit et al. \(2005\)](#) or [Föllmer and Schied \(2016\)](#)). As for comonotonic subadditivity, less attention has been devoted to this property. [Song and Yan \(2006\)](#) provided, along with further properties, a full characterization of comonotonic subadditive functionals as envelopes of Choquet integrals, and [Song and Yan \(2009\)](#) provided some applications thereof.

Remark 2. Proposition 1 hints towards the possibility of representing continuous, monotone, positively homogeneous, and strictly comonotonic subadditive functionals as suprema of signed Choquet integrals. Indeed, as shown by [Wang et al. \(2020\)](#), comonotonic additive and continuous functionals can be represented by Choquet integrals with respect to signed capacities.

6. CONCLUDING REMARKS AND AN OPEN QUESTION

Building upon the work of [Amarante \(2019\)](#) and [Fuchssteiner and Wright \(1977\)](#), we provided a nonlinear version of the classical Sandwich Theorem. We used this result to retrieve an extension result and an envelope representation result. Our examples show that the type of nonlinearities that we introduce include some important cases that have been devoted considerable attention in decision theory and mathematical finance. Our approach highlights the possibility of retrieving Hahn-Banach-type extension results that are widely applied in functional analysis, theoretical economics, and mathematical finance. We conclude with an (informal) question and two conjectures:

- (O1) To what extent do our results depend on the fact that our maps can take $-\infty$ as a value? More formally, is it possible to prove the following reformulations of Corollaries 1 and 2?

Conjecture 1. *Let $H : X \rightarrow \mathbb{V}$ be \mathbf{C} -sublinear and monotone, and let $Y \subseteq X$ be a convex cone. If $\ell : Y \rightarrow \mathbb{V}$ is \mathbf{C} -linear and satisfies $\ell \leq H|_Y$, then there exists a \mathbf{C} -linear $Q : X \rightarrow \mathbb{V}$ such that $Q \leq H$ and $\ell \leq Q|_Y$.*

Conjecture 2. *If $H : X \rightarrow \mathbb{V}$ is \mathbf{C} -sublinear and monotone, then*

$$H(x) = \sup_{Q \in \tilde{D}(H)} Q(x), \text{ for all } x \in X,$$

where

$$\tilde{D}(H) = \left\{ Q : X \rightarrow \mathbb{V} \mid Q \text{ is } \mathbf{C}\text{-linear and } Q \leq H \right\}.$$

APPENDIX A. ADDITIONAL RESULTS FOR SECTION 5

In this section we report some auxiliary results that we applied in Section 5. In particular, we provide a partial answer to the following

Question 1. If $x, y \in B(\Omega, \mathcal{F})$ are comonotonic, then there exist two sequences $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}$ in $B(\Omega, \mathcal{F})$ such that:

- (1) $x_n \rightarrow x$ and $y_n \rightarrow y$;
- (2) for all $n \in \mathbb{N}$, x_n and y_n are strictly comonotonic.

It is immediate to see that this does not hold in general measurable spaces. Indeed, take any $\Omega \neq \emptyset$ and let $\mathcal{F} = \{\emptyset, \Omega\}$. In this measurable space (Ω, \mathcal{F}) , a function is \mathcal{F} -measurable if and only if it is constant, and therefore there is no injective measurable function. This observation highlights the fact that Question 1 may admit a positive answer only if we focus on measurable spaces with a sufficiently sparse sigma-algebra, where this sparsity depends also on the cardinality of Ω . Providing a full answer to this question is out of the scope of this paper. Hence, we focus on a special case. However, we first need some auxiliary lemmas.

Lemma 5. *Let $x, y \in B(\Omega, \mathcal{F})$. The following are equivalent*

- (i) *x and y are strictly comonotonic with $x \notin C_y$.*
- (ii) *there exist two increasing functions $h, g : \mathbb{R} \rightarrow \mathbb{R}$ and an injective $z \in B(\Omega, \mathcal{F})$ such that $x = h(z)$, $y = g(z)$, and h, g are injective over $z(\Omega)$.*

Proof. (i) \Rightarrow (ii). Since x, y are comonotonic, there exist increasing functions $h, g : \mathbb{R} \rightarrow \mathbb{R}$ and $z \in B(\Omega, \mathcal{F})$ such that $x = h(z)$ and $y = g(z)$ (see e.g., [Denuit et al. \(2023\)](#), Theorem 2.7). Suppose that z is not injective. Then, there exist $\omega_1, \omega_2 \in \Omega$ such that $\omega_1 \neq \omega_2$ and $z(\omega_1) = z(\omega_2)$. Thus,

$$[x(\omega_1) - x(\omega_2)] [y(\omega_1) - y(\omega_2)] = [h(z(\omega_1)) - h(z(\omega_2))] [y(\omega_1) - y(\omega_2)] = 0,$$

contradicting the strict comonotonicity of x, y . Therefore, z must be injective. Now suppose that h is not injective over $z(\Omega)$. Then there exist $\omega_1, \omega_2 \in \Omega$ such that $\omega_1 \neq \omega_2$, $z(\omega_1) \neq z(\omega_2)$, and $h(z(\omega_1)) = h(z(\omega_2))$. Since z is injective, we have that $\omega_1 \neq \omega_2$ and

$$[x(\omega_1) - x(\omega_2)] [y(\omega_1) - y(\omega_2)] = [h(z(\omega_1)) - h(z(\omega_2))] [y(\omega_1) - y(\omega_2)] = 0,$$

contradicting the strict comonotonicity of x, y . Interchanging h with g and x with y , the same conclusion holds for g as well.

(ii) \Rightarrow (i). If $\omega_1 \neq \omega_2$ and $h(z(\omega_1)) > h(z(\omega_2))$, then we must have that $z(\omega_1) > z(\omega_2)$ and hence $g(z(\omega_1)) > g(z(\omega_2))$. This proves the claim. \square

Lemma 6. *Suppose that $f : [a, b] \rightarrow \mathbb{R}$ is 1-Lipschitz and increasing. Then, there exists a sequence of 1-Lipschitz and strictly increasing functions $(f_n)_{n \in \mathbb{N}}$ from $[a, b]$ to \mathbb{R} that converges uniformly to f .*

Proof. Since f is continuous and increasing, there exist at most countably many disjoint nondegenerate intervals $(I_n)_{n \in \mathbb{N}}$ over which f is constant, i.e., $f(I_n) = \{k_n\}$ for all $n \in \mathbb{N}$ and some $k_n \in \mathbb{R}$. Fix $\varepsilon > 0$ and take finitely many points, $a = a_1 < \dots < a_k = b$ such that $|a_i - a_{i+1}| < \varepsilon/2$ for all $i = 1, \dots, k-1$. Suppose that $f(a_i) = f(a_j)$ for some $i < j$. Since f is increasing, we have that

$$f(a_i) = f(a_{i+1}) = \dots = f(a_j).$$

This implies that $a_i, \dots, a_j \in I_n$ for some $n \in \mathbb{N}$. Thus, to find a strictly increasing approximation we need to modify our vector of points. In particular, we remove all points a_i, \dots, a_{j-1} , and we add one point \tilde{a}_i picked from the set

$$\left\{ x \in [a_{i-1}, \inf I_n) : f(x) < f(a_i) \text{ and } |f(x) - f(a_i)| < \frac{\varepsilon}{2} \right\}.$$

Repeating this operation for all points where the function is constant, within the set $\{a_1, \dots, a_k\}$, we retrieve (in a finite amount of operations) a set $\{\tilde{a}_1, \dots, \tilde{a}_m\}$ with $\tilde{a}_1 = a$ and $\tilde{a}_m = b$ such that

$$f(\tilde{a}_1) < f(\tilde{a}_2) < \dots < f(\tilde{a}_{m-1}) < f(\tilde{a}_m).$$

Define the function $f^m : \{\tilde{a}_1, \dots, \tilde{a}_m\} \rightarrow \mathbb{R}$ by $f^m(\tilde{a}_i) = f(\tilde{a}_i)$, for all $i = 1, \dots, m$. By linear interpolation, we can extend f^m to the whole interval $[a, b]$, and we denote such an extension again by f^m . Clearly, f^m is strictly increasing, and we now show that it must also be 1-Lipschitz. To this end, notice that the slope of $f^m|_{[\tilde{a}_i, \tilde{a}_{i+1}]}$ satisfies

$$\frac{f(\tilde{a}_{i+1}) - f(\tilde{a}_i)}{\tilde{a}_{i+1} - \tilde{a}_i} \leq 1,$$

for all $i = 1, \dots, k-1$, where the inequality follows from the fact that f is 1-Lipschitz, $\tilde{a}_{i+1} > \tilde{a}_i$, and f is increasing. This implies that $f^m|_{[\tilde{a}_i, \tilde{a}_{i+1}]}$ is 1-Lipschitz for all $i = 1, \dots, k-1$. If $x > y$, then $x \in (\tilde{a}_i, \tilde{a}_{i+1}]$ and $y \in [\tilde{a}_j, \tilde{a}_{j+1}]$ for some $i > j$. This implies that

$$\begin{aligned} |f^m(x) - f^m(y)| &= f^m(x) - f^m(y) \\ &= f^m(x) - f^m(\tilde{a}_i) + f^m(\tilde{a}_i) - f^m(\tilde{a}_{i-1}) + \dots + f^m(\tilde{a}_{j+1}) - f^m(\tilde{a}_j) + f^m(\tilde{a}_j) - f^m(y) \\ &\leq x - \tilde{a}_i + \tilde{a}_i - \tilde{a}_{i-1} + \dots + \tilde{a}_{j+1} - \tilde{a}_j + \tilde{a}_j - y = |x - y|. \end{aligned}$$

Thus f^m is 1-Lipschitz. Now we prove that f^m is ε -close to f . For all $x \in [\tilde{a}_{i-1}, \tilde{a}_i]$ and $i = 2, \dots, m$ we have

$$\begin{aligned} |f(x) - f^m(x)| &\leq |f(x) - f(\tilde{a}_{i-1})| + |f(\tilde{a}_{i-1}) - f^m(x)| \\ &\leq \frac{\varepsilon}{2} + |f^m(\tilde{a}_{i-1}) - f^m(\tilde{a}_i)| \\ &= \frac{\varepsilon}{2} + |f(\tilde{a}_{i-1}) - f(\tilde{a}_i)| \\ &< \varepsilon. \end{aligned}$$

Since x was chosen arbitrarily, it follows that $\|f - f^m\|_\infty \rightarrow 0$. Therefore, there exists a sequence of 1-Lipschitz and strictly increasing functions $(f_n)_{n \in \mathbb{N}}$ converging uniformly to f . \square

Now suppose that $\Omega = [0, 1]$ and $\mathcal{F} = \mathcal{B}[0, 1]$ is the Borel sigma-algebra.⁵

Lemma 7. *If $x, y \in B(\Omega, \mathcal{F})$ are comonotonic, then there exist two sequences $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}$ in $B(\Omega, \mathcal{F})$ such that:*

- (1) $x_n \xrightarrow{\|\cdot\|_\infty} x$ and $y_n \xrightarrow{\|\cdot\|_\infty} y$;
- (2) For all $n \in \mathbb{N}$, x_n and y_n are strictly comonotonic with $x_n \notin C_{y_n}$.

Proof. Since $x, y \in B(\Omega, \mathcal{F})$ are comonotonic, there exist two increasing 1-Lipschitz functions $h, g : \mathbb{R} \rightarrow \mathbb{R}$ and an $z \in B(\Omega, \mathcal{F})$ such that $x = h(z)$, $y = g(z)$ (Denuit et al. (2023), Theorem 2.7). We first prove a claim that will yield the result.

CLAIM. There exists a sequence of injective and measurable functions converging to z .

Proof of the claim. Let $(s_n)_{n \in \mathbb{N}} \in B(\Omega, \mathcal{F})^{\mathbb{N}}$ be a sequence of step functions converging uniformly to z . Each s_n can be uniquely identified with a partition $(I_i^n)_{i=1}^{k_n}$ of nondegenerate subintervals of Ω , and a vector of values $(a_1^n, \dots, a_{k_n}^n)$, for some $k_n \in \mathbb{N}$. For all $n \in \mathbb{N}$ and $\varepsilon > 0$, we can define the following,

$$s_n^\varepsilon(\omega) = 2\varepsilon \left(\frac{\omega - \inf I_i^n}{\sup I_i^n - \inf I_i^n} \right) + a_i^n - \varepsilon,$$

for all $\omega \in I_i^n$ and all $i = 1, \dots, k_n$. Clearly s_n^ε is an injective Borel measurable function, for all $\varepsilon > 0$ and all $n \in \mathbb{N}$. Intuitively, we are simply rotating slightly the constant “lines” of each s_n over all their partitions. Moreover, note that

$$\|s_n^\varepsilon - s_n\|_\infty \leq \varepsilon,$$

for all $n \in \mathbb{N}$ and $\varepsilon > 0$. This implies that $(s_n^{1/n})_{n \in \mathbb{N}}$ converges uniformly to z . Indeed,

$$\left\| z - s_n^{1/n} \right\|_\infty \leq \|z - s_n\|_\infty + \left\| s_n - s_n^{1/n} \right\|_\infty \leq \|z - s_n\|_\infty + \frac{1}{n} \rightarrow 0.$$

Thus, we found a sequence of injective and measurable functions converging uniformly to z . \square

Given that z is bounded, there exist $m, M \in \mathbb{R}$ such that $z(\Omega) \subseteq (m, M)$. Since $(s_n^{1/n})_{n \in \mathbb{N}}$ converges uniformly to z , there exists some $N \in \mathbb{N}$ sufficiently large so that

$$s_n^{1/n}(\Omega) \subseteq [m, M],$$

for all $n \geq N$. Using a slight abuse of notation, we will now identify by $(s_n^{1/n})_{n \in \mathbb{N}}$ its subsequence $(s_{n_k}^{1/n_k})_{k \in \mathbb{N}}$ with $n_1 = N$, $n_2 = N + 1$ and so on. By Lemma 6, there exist two sequences of 1-Lipschitz and strictly increasing functions $(h_n)_{n \in \mathbb{N}}$ and $(g_n)_{n \in \mathbb{N}}$ from $[m, M]$ to

⁵All the results provided in this section would hold for any closed interval $I \subseteq \mathbb{R}$.

\mathbb{R} converging uniformly to h and g . For all $n \in \mathbb{N}$, let $x_n = h_n \left(s_n^{1/n} \right)$ and $y_n = g_n \left(s_n^{1/n} \right)$. Fix $n \in \mathbb{N}$ arbitrarily. If $\omega_1 \neq \omega_2$, then $s_n^{1/n}(\omega_1) \neq s_n^{1/n}(\omega_2)$ since $s_n^{1/n}$ is injective, say without loss of generality that $s_n^{1/n}(\omega_1) > s_n^{1/n}(\omega_2)$. Since h_n and g_n are both strictly increasing we have that

$$\begin{aligned} & [x_n(\omega_1) - x_n(\omega_2)] [y_n(\omega_1) - y_n(\omega_2)] \\ &= \left[h_n \left(s_n^{1/n} \right) (\omega_1) - h_n \left(s_n^{1/n} \right) (\omega_2) \right] \left[g_n \left(s_n^{1/n} \right) (\omega_1) - g_n \left(s_n^{1/n} \right) (\omega_2) \right] > 0. \end{aligned}$$

Thus, x_n, y_n are strictly comonotonic. Fix $n \in \mathbb{N}$ arbitrarily. Since h_n is 1-Lipschitz, we have

$$\begin{aligned} \left\| h(z) - h_n \left(s_n^{1/n} \right) \right\|_{\infty} &\leq \|h(z) - h_n(z)\|_{\infty} + \left\| h_n(z) - h_n \left(s_n^{1/n} \right) \right\|_{\infty} \\ &= \|h(z) - h_n(z)\|_{\infty} + \sup_{\omega \in \Omega} \left| h_n(z(\omega)) - h_n \left(s_n^{1/n}(\omega) \right) \right| \\ &\leq \|h(z) - h_n(z)\|_{\infty} + \sup_{\omega \in \Omega} \left| z(\omega) - s_n^{1/n}(\omega) \right| \rightarrow 0. \end{aligned}$$

Thus $(x_n)_{n \in \mathbb{N}}$ converges uniformly to x . The same holds for $(y_n)_{n \in \mathbb{N}}$, and the proof is totally analogous. \square

REFERENCES

- Aliprantis, C. D. and Border, K. C. (2006). *Infinite dimensional analysis*. Springer, Berlin, third edition. A hitchhiker's guide.
- Aliprantis, C. D. and Burkinshaw, O. (2006). *Positive operators*. Springer, Dordrecht. Reprint of the 1985 original.
- Amarante, M. (2019). The sandwich theorem via Pataia's fixed point theorem. *Positivity*, 23(1):97–100.
- Caspar, N., Leclerc, B., and Monjardet, B. (2012). *Finite ordered sets*, volume 144 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge. Concepts, results and uses.
- Denuit, M., Dhaene, J., Ghossoub, M., and Robert, C. Y. (2023). Comonotonicity and pareto optimality, with application to collaborative insurance. *Mimeo*.
- Denuit, M., Dhaene, J., Goovaerts, M., and Kaas, R. (2005). *Actuarial Theory for Dependent Risks: Measures, Orders and Models*. John Wiley & Sons.
- Escardó, M. H. (2003). Joins in the frame of nuclei. *Appl. Categ. Structures*, 11(2):117–124.
- Föllmer, H. and Schied, A. (2016). *Stochastic Finance: An Introduction in Discrete Time – 4th ed.* Walter de Gruyter.
- Fuchssteiner, B. and Wright, J. D. M. (1977). Representing isotone operators on cones. *Quart. J. Math. Oxford Ser. (2)*, 28(110):155–162.
- König, H. (1972). Sublineare Funktionale. *Arch. Math. (Basel)*, 23:500–508.

- Pataraiia, D. (1997). A constructive proof of tarski's fixed-point theorem for dcpos. In *65th Peripatetic Seminar on Sheaves and Logic*.
- Roth, W. (2000). Hahn-Banach type theorems for locally convex cones. *J. Austral. Math. Soc. Ser. A*, 68(1):104–125.
- Schmeidler, D. (1986). Integral representation without additivity. *Proc. Amer. Math. Soc.*, 97(2):255–261.
- Schmeidler, D. (1989). Subjective probability and expected utility without additivity. *Econometrica*, 57(3):571–587.
- Schröder, B. (2016). *Ordered sets*. Birkhäuser/Springer, second edition. An introduction with connections from combinatorics to topology.
- Song, Y. and Yan, J. (2006). The representations of two types of functionals on $L^\infty(\Omega, \mathcal{F})$ and $L^\infty(\Omega, \mathcal{F}, \mathbb{P})$. *Sci. China Ser. A*, 49(10):1376–1382.
- Song, Y. and Yan, J.-A. (2009). Risk measures with comonotonic subadditivity or convexity and respecting stochastic orders. *Insurance Math. Econom.*, 45(3):459–465.
- Wang, R., Wei, Y., and Willmot, G. E. (2020). Characterization, robustness, and aggregation of signed Choquet integrals. *Math. Oper. Res.*, 45(3):993–1015.

Mario Ghossoub: UNIVERSITY OF WATERLOO – DEPARTMENT OF STATISTICS AND ACTUARIAL SCIENCE
– 200 UNIVERSITY AVE. W. – WATERLOO, ON, N2L 3G1 – CANADA

Email address: mario.ghossoub@uwaterloo.ca

Giulio Principi: NEW YORK UNIVERSITY – DEPARTMENT OF ECONOMICS – 19 WEST 4TH STREET –
NEW YORK, NY 10012 – USA

Email address: gp2187@nyu.edu

Lorenzo Stanca: COLLEGIO CARLO ALBERTO AND UNIVERSITY OF TURIN – DEPARTMENT OF ESOMAS
– CORSO UNIONE SOVIETICA, 218 BIS – TURIN 10134 – ITALY

Email address: lorenzo.stanca@carloalberto.org